

SCHEDULING STATUS:

S4

ADCO ACYCLOVIR 200 mg tablets

Acyclovir

Contains sugar (lactose): 170 mg per tablet

Read all of this leaflet carefully before you start taking ADCO ACYCLOVIR 200 mg

- Keep this leaflet. You may need to read it again.
- If you have further questions, please ask your doctor, pharmacist, nurse, or other healthcare provider.
- ADCO ACYCLOVIR 200 mg has been prescribed for you personally and you should not share your medicine with other people. It may harm them, even if their symptoms are the same as yours.

What is in this leaflet

1. What ADCO ACYCLOVIR 200 mg is and what it is used for
2. What you need to know before you take ADCO ACYCLOVIR 200 mg
3. How to take ADCO ACYCLOVIR 200 mg
4. Possible side effects
5. How to store ADCO ACYCLOVIR 200 mg
6. Contents of the pack and other information

1. What ADCO ACYCLOVIR 200 mg is and what it is used for

ADCO ACYCLOVIR 200 mg belongs to group of medicines called antivirals. ADCO ACYCLOVIR 200 mg is used for the treatment of the following:

- Herpes simplex infections of the skin and mucous membranes including initial and recurrent genital Herpes simplex virus infections.
- Suppression of recurrent Herpes simplex infections in immunocompetent patients.
- Prophylaxis of Herpes simplex infections in immunocompromised patients.
- Treatment of Herpes zoster if the lesions are not older than 72 hours.
- Treatment of Varicella zoster (chickenpox) within 24 hours after appearance of the typical chickenpox rash.

2. What you need to know before you take ADCO ACYCLOVIR 200 mg

Do not take ADCO ACYCLOVIR 200 mg:

- If you are hypersensitive (allergic) to acyclovir, valaciclovir or any of the other ingredients of ADCO ACYCLOVIR 200 mg (listed in section 6).
- If you are pregnant or breastfeeding your baby

Warnings and precautions

Take special care with ADCO ACYCLOVIR 200 mg:

- If you are elderly or have kidney problems, you may have a higher risk of side-effects and your doctor may watch you more closely.
- If you have kidney problems. Your doctor may prescribe a lower dose of ADCO ACYCLOVIR 200 mg.
- It is important that you drink plenty of water while taking high doses of ADCO ACYCLOVIR 200 mg.
- Serious skin reactions such as Drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported in patients receiving ADCO ACYCLOVIR 200 mg. Combination of symptoms such as fever, severe rash, peeling skin, swelling of the face, swollen lymph glands, flu-like feeling, yellow skin or eyes, shortness of breath, dry cough, chest pain or discomfort, feel thirsty, urinating less often, less urine can be seen. If you experience any signs of serious skin reactions such as swelling, itching, red severe rash, stop using ADCO ACYCLOVIR 200 mg immediately and contact your doctor (see Section 4).

Other medicines and ADCO ACYCLOVIR 200 mg

Always tell your healthcare provider if you are taking any other medicine. (This includes all complementary or traditional medicines.)

In particular tell your doctor or pharmacist if you are taking any of the following medicines:

- probenecid, used to treat gout
- cimetidine, used to treat stomach ulcers
- mycophenolate mofetil, used to stop your body rejecting transplanted organs.
- theophylline, used to treat asthma.

Pregnancy, breastfeeding and fertility

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other healthcare provider for advice before using ADCO ACYCLOVIR 200 mg.

Driving and using machines

ADCO ACYCLOVIR 200 mg can cause dizziness. You should not drive or operate machinery if you experience this effect.

It is not always possible to predict to what extent ADCO ACYCLOVIR 200 mg may interfere with the daily activities of a patient. Patients should ensure that they do not engage in the above activities until they are aware of the measure to which ADCO ACYCLOVIR 200 mg affects them.

ADCO ACYCLOVIR 200 mg contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact

your doctor before taking ADCO ACYCLOVIR 200 mg.

ADCO ACYCLOVIR 200 mg contains sodium

ADCO ACYCLOVIR 200 mg contains less than 1mmol sodium (23 mg) per dosage unit, that is to say essentially 'sodium-free'.

3. How to take ADCO ACYCLOVIR 200 mg

Do not share medicines prescribed for you with any other person.

Always use ADCO ACYCLOVIR 200 mg exactly as your doctor or pharmacist has told you.

Check with your doctor or pharmacist if you are not sure.

Your doctor will tell you how long your treatment with ADCO ACYCLOVIR 200 mg will last.

If you have the impression that the effect of ADCO ACYCLOVIR 200 mg is too strong or too weak, tell your doctor or pharmacist.

Dosage in adults:

Initial and recurrent herpes simplex infections of skin and mucous membranes:

200 mg (one ADCO Acyclovir 200 mg tablet) 5 times daily at 4 hourly intervals for 5 days, omitting the night dose. In severe initial infection, your doctor will extend your treatment. In immunocompromised patients, dosage could be increased to 400 mg (two ADCO ACYCLOVIR 200 mg tablets), or alternatively, intravenous dosing can be considered

The first dose should be administered as early as possible after start of infection

Suppression of recurrent genital herpes simplex infections in patients with a weakened immune system:

200 mg acyclovir (one ADCO ACYCLOVIR 200 mg tablet) 4 times daily at 6 hourly intervals, or 400 mg (two ADCO ACYCLOVIR 200 mg tablets) 12 hourly.

The dose may be reduced to 200 mg acyclovir (one ADCO ACYCLOVIR 200 mg tablet) every 8 hours or even every 12 hours.

Some patients may have breakthrough infections on 800 mg (four ADCO ACYCLOVIR 200 mg tablets) daily. Your doctor will interrupt therapy every 6 – 12 months to observe history of disease.

Prevention of herpes simplex infections in patients with a weakened immune system:

200 mg 4 times daily at 6 hourly intervals.

In severely immunocompromised patients, or impaired absorption from the gut, dosage could be doubled to 400 mg (two ADCO ACYCLOVIR 200 mg tablets) or alternatively, intravenous dosing can be considered.

Treatment of varicella zoster in adolescents (12-18 years):

A dose of 800 mg (four ADCO ACYCLOVIR 200 mg tablets) should be taken daily for 5 days.

Treatment of varicella zoster and herpes zoster infections in adults:

800 mg (four ADCO ACYCLOVIR 200 mg tablets) 5 times daily at 4 hourly intervals for 7 days, omitting the night dose

Dosage in the elderly:

Adequate hydration in patients taking high doses must be maintained. Patients with kidney problems should be given special attention.

Treatment or prophylaxis of herpes simplex infections in patients with renal impairment:

200 mg every 12 hours if creatinine clearance 0-10 mL/minute

Treatment of varicella zoster in immunocompromised patients with renal impairment:
800 mg (four ADCO ACYCLOVIR 200 mg tablets) every 12 hours if creatinine clearance 0-10 mL/minute

800 mg (four ADCO ACYCLOVIR 200 mg tablets) every 8 hourly intervals if creatinine clearance 10-25 mL/minute

800 mg (four ADCO ACYCLOVIR 200 mg tablets) 5 times daily with 4 hourly intervals if creatinine clearance > 25 mL/minute.

Patients on haemodialysis should receive their usual appropriate dose after each dialysis

Orally administered acyclovir in children less than 2 years of age has not yet been fully studied

The tablets should be taken by mouth

If you take more ADCO ACYCLOVIR 200 mg than you should

In the event of overdosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison centre.

If you forget to take ADCO ACYCLOVIR 200 mg

Take it as soon as you remember. However, if it is nearly time for your next dose, skip the missed dose.

Do not take a double dose to make up for a forgotten dose.

4. Possible side effects

ADCO ACYCLOVIR 200 mg can have side effects.

Not all side effects reported for ADCO ACYCLOVIR 200 mg are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking ADCO ACYCLOVIR 200 mg, please consult your health care provider for advice.

If any of the following happens, stop taking ADCO ACYCLOVIR 200 mg and tell your doctor immediately or go to the casualty department at your nearest hospital:

Frequent side effects include:

- headache
- feeling dizzy
- nausea
- diarrhoea
- vomiting
- stomach pains
- itchy skin (pruritis)
- skin rash
- skin reaction after exposure to light (photosensitivity)
- hives (urticaria)
- hair loss
- feeling tired
- fever

Less frequent side effects include:

- effects on some blood tests
- anaphylaxis (life-threatening allergic reaction)

Angioedema (rapid swelling under the skin)

- agitation, confusion, shakiness, impaired coordination, slurred speech, hallucinations, psychotic symptoms, fits, drowsiness, brain diseases, coma
- shortness of breath
- Reversible rises in bilirubin and liver related enzymes
- inflammation of the liver (hepatitis)
- yellowing of your skin and whites of your eyes (jaundice)
- Increased blood urea and creatinine
- renal failure, renal pain

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects

If you get side effects, talk to your doctor, pharmacist or nurse. You can also report side effects to SAHPRA via the “**6.04 Adverse Drug Reaction Reporting Form**”, found online under SAHPRA’s publications: <https://www.sahpra.org.za/Publications/Index/8>. By reporting side effects, you can help provide more information on the safety of ADCO ACYCLOVIR 200 mg.

You may also report to Adcock Ingram Limited using the following e-mail address: Adcock.AEReports@adcock.com.

5. How to store ADCO ACYCLOVIR 200 mg

- Store all medicines out of reach of children.

- Store at or below 25°C in dry place. Protect from light.
- Store in the original package.
- Do not use after the expiry date stated on the label / carton / bottle
- Return all unused medicine to your pharmacist.
- Do not dispose of unused medicine in drains or sewerage systems (e.g., toilets).

6. Contents of the pack and other information

What ADCO ACYCLOVIR 200 mg contains

The active ingredient is acyclovir 200 mg.

The other ingredients are lactose, colloidal silicon dioxide, magnesium stearate, microcrystalline cellulose, pregelatinized maize starch, sodium lauryl sulphate, sodium starch glycollate and stearic acid.

What ADCO ACYCLOVIR 200 mg looks like and contents of the pack

ADCO ACYCLOVIR 200 mg are: White tablets, hexagonal in shape, scored line on one side.

ADCO ACYCLOVIR 200 mg are available in:

Pack size of 25 tablets packed in a white polypropylene securitainer with a white LDPE closure.

Pack size of 25 tablets packed in blisters using PVC film and aluminium foil.

Not all pack sizes may be marketed.

Holder of Certificate of Registration

Adcock Ingram Limited

1 New Road,

Erand Gardens,

Midrand,

1685

Customer Care: 0860 ADCOCK / 232625

This leaflet was last revised in

08 November 2024

Registration number

28/20.2.8/0552

Botswana (S2): BOT1803283
Namibia (NS2): 04/20.2.8/1554

SKEDULERINGSSTATUS: S4

ADCO ACYCLOVIR 200 mg tablette
Asiklovir
Bevat suiker (laktose): 170 mg per tablet

Lees hierdie hele biljet noukeurig deur voordat u ADCO ACYCLOVIR 200 mg begin neem

- Hou hierdie biljet. U sal dit dalk weer moet lees.
- Indien u verdere vrae het, raadpleeg asseblief u dokter, apteker, verpleegkundige of ander gesondheidsorgverskaffer.
- ADCO ACYCLOVIR 200 mg is aan u persoonlik voorgeskryf en u moenie u medisyne met ander mense deel nie. Dit kan hulle benadeel, selfs al ervaar hul dieselfde simptome as u.

Wat is in hierdie biljet

1. Wat ADCO ACYCLOVIR 200 mg is en waarvoor dit gebruik word
2. Wat u moet weet voordat u ADCO ACYCLOVIR 200 mg neem
3. Hoe om ADCO ACYCLOVIR 200 mg te neem
4. Moontlike nuwe-effekte
5. Hoe om ADCO ACYCLOVIR 200 mg te bêre
6. Inhoud van die verpakking en ander inligting

1. Wat ADCO ACYCLOVIR 200 mg is en waarvoor dit gebruik word

ADCO ACYCLOVIR 200 mg behoort aan die groep medisyne wat antivirale middels genoem word. ADCO ACYCLOVIR 200 mg word gebruik vir die behandeling van die volgende:

- *Herpes simplex* infeksies van die vel en slymvliese, insluitend aanvanklike en herhalende genitale *Herpes simplex*-virus infeksies.
- Onderdrukking van herhalende *Herpes simplex* infeksies by immuunbevoegde pasiënte.
- Profilakse van *Herpes simplex* infeksies by immuunonderdrukte pasiënte.
- Behandeling van *Herpes zoster*, indien die letsels nie ouer as 72 uur is nie.
- Behandeling van *Varicella zoster* (waterpokkies) binne 24 uur ná die voorkoms van die tipiese waterpokkie-uitslag.

2. Wat u moet weet voordat u ADCO ACYCLOVIR 200 mg neem

Moenie ADCO ACYCLOVIR 200 mg neem nie:

- Indien u hipersensitief (allergies) is vir asiklovir, valasiklovir of enige van die ander bestanddele van ADCO ACYCLOVIR 200 mg (gelys in afdeling 6).
- Indien u swanger is of u baba borsvoed

Waarskuwings en voorsorgmaatreëls

Wees veral versigtig met ADCO ACYCLOVIR 200 mg:

- Indien u bejaard is of nierprobleme het, kan u 'n groter risiko vir newe-effekte hê en u dokter kan u meer noukeurig dophou.
- Indien u nierprobleme het. U dokter kan 'n laer dosis ADCO ACYCLOVIR 200 mg voorskryf.
- Dit is belangrik dat u baie water drink terwyl u hoë dosisse ADCO ACYCLOVIR 200 mg neem.
- Ernstige velreaksies soos Geneesmiddelreaksie met eosinofilie en sistemiese simptome (GRESS) is aangemeld by pasiënte wat ADCO ACYCLOVIR 200 mg ontvang. 'n Kombinasie van simptome soos koors, erge veluitslag, afskilferende vel, swelling van die gesig, geswelde limfkliere, griepagtige gevoel, geel vel of oë, kortasem, droë hoes, borspyn of -ongemak, voel dors, urineer minder gereeld en minder urien word opgemerk. Indien u enige tekens van ernstige velreaksies soos swelling, jeukerigheid, rooi ernstige veluitslag ervaar, staak die gebruik van ADCO ACYCLOVIR 200 mg onmiddellik en kontak u dokter (sien Afdeling 4).

Ander medisyne en ADCO ACYCLOVIR 200 mg

Lig altyd u gesondheidsorgverskaffer in indien u enige ander medisyne neem. (Dit sluit alle komplementêre of tradisionele medisyne in.)

Lig veral u dokter of apteker in indien u enige van die volgende medisyne neem:

- probenesied, wat gebruik word om jig te behandel
- simetidien, wat gebruik word om maagsere te behandel
- mikofenolaatmofetiel, wat gebruik word om te keer dat u liggaam oorgeplante organe verwerp.
- teofillien, wat gebruik word om asma te behandel.

Swangerskap, borsvoeding en vrugbaarheid

Indien u swanger is of borsvoed, dink u is dalk swanger of van plan is om 'n baba te hê, raadpleeg asseblief u dokter, apteker of ander gesondheidsorgverskaffer vir advies voordat u ADCO ACYCLOVIR 200 mg gebruik.

Bestuur en gebruik van masjinerie

ADCO ACYCLOVIR 200 mg kan duiseligheid veroorsaak. U moet nie bestuur of masjinerie hanteer indien u hierdie effek ervaar nie.

Dit is nie altyd moontlik om te voorspel in watter mate ADCO ACYCLOVIR 200 mg met die daaglikse aktiwiteite van 'n pasiënt kan inmeng nie. Pasiënte moet verseker dat hulle nie bogenoemde aktiwiteite uitvoer totdat hulle bewus is van die mate waartoe ADCO ACYCLOVIR 200 mg hulle beïnvloed nie.

ADCO ACYCLOVIR 200 mg bevat laktose

Indien u deur u dokter ingelig is dat u 'n onverdraagsaamheid teenoor sommige suikers het,

kontak u dokter voordat u ADCO ACYCLOVIR 200 mg neem.

ADCO ACYCLOVIR 200 mg bevat natrium

ADCO ACYCLOVIR 200 mg bevat minder as 1 mmol natrium (23 mg) per dosis-eenheid, dit wil sê in wese 'natriumvry'.

3. Hoe om ADCO ACYCLOVIR 200 mg te neem

Moenie medisyne wat vir u voorgeskryf is, met enige ander persoon deel nie.

Gebruik altyd ADCO ACYCLOVIR 200 mg presies soos u dokter of apteker u ingelig het.

Raadpleeg u dokter of apteker indien u nie seker is nie.

U dokter sal u inlig hoe lank u behandeling met ADCO ACYCLOVIR 200 mg sal duur.

Indien u die indruk kry dat die effek van ADCO ACYCLOVIR 200 mg te sterk of te swak is, raadpleeg u dokter of apteker.

Dosis by volwassenes:

Aanvanklike en herhalende *herpes simplex* infeksies van vel en slymvliese:

200 mg (een ADCO ACYCLOVIR 200 mg tablet) 5 keer per dag met 4-uurlikse tussenposes vir 5 dae, met weglating van die nagdosis. In ernstige aanvanklike infeksie sal u dokter u behandeling verleng. By immuunonderdrukte pasiënte kan die dosis verhoog word tot 400 mg (twee ADCO ACYCLOVIR 200 mg tablette), of alternatiewelik kan intraveneuse dosering oorweeg word.

Die eerste dosis moet so vroeg as moontlik ná die aanvang van die infeksie toegedien word.

Onderdrukking van herhalende genitale *herpes simplex* infeksies by pasiënte met 'n verswakte immuunstelsel:

200 mg asiklovir (een ADCO ACYCLOVIR 200 mg tablet) 4 keer per dag met 6-uurlikse tussenposes, of 400 mg (twee ADCO ACYCLOVIR 200 mg tablette) 12-uurliks.

Die dosis kan verminder word tot 200 mg asiklovir (een ADCO ACYCLOVIR 200 mg tablet) elke 8 uur of selfs elke 12 uur.

Sommige pasiënte kan deurbraak-infeksies op 800 mg (vier ADCO ACYCLOVIR 200 mg tablette) daagliks hê. U dokter sal elke 6–12 maande die terapie onderbreek om die geskiedenis van die siekte waar te neem.

Voorkoming van *herpes simplex* infeksies by pasiënte met 'n verswakte immuunstelsel:

200 mg 4 keer per dag met 6-uurlikse tussenposes.

By pasiënte met ernstige immuunonderdrukking, of verswakte absorpsie uit die ingewande, kan die dosis verdubbel word tot 400 mg (twee ADCO ACYCLOVIR 200 mg tablette), of alternatiewelik kan intraveneuse dosering oorweeg word.

Behandeling van *varicella zoster* by adolessente (12-18 jaar):

'n Dosis van 800 mg (vier ADCO ACYCLOVIR 200 mg tablette) moet daagliks vir 5 dae

geneem word.

Behandeling van *varicella zoster* en *herpes zoster* infeksies by volwassenes:

800 mg (vier ADCO ACYCLOVIR 200 mg tablette) 5 keer per dag met 4-uurlikse tussenposes vir 7 dae, met weglating van die nagdosis.

Dosis by bejaardes:

By pasiënte wat hoë dosisse neem, moet voldoende hidrasie gehandhaaf word. Pasiënte met nierprobleme moet spesiale aandag geniet.

Behandeling of voorkoming van *herpes simplex* infeksies by pasiënte met nierinkorting:

200 mg elke 12 uur indien kreatinienopruiming 0-10 mL/minuut is.

Behandeling van *varicella zoster* by immuunonderdrukte pasiënte met nierinkorting:

800 mg (vier ADCO ACYCLOVIR 200 mg tablette) elke 12 uur indien kreatinienopruiming 0-10 mL/minuut is.

800 mg (vier ADCO ACYCLOVIR 200 mg tablette) met 8-uurlikse intervale indien kreatinienopruiming 10-25 mL/minuut is.

800 mg (vier ADCO ACYCLOVIR 200 mg tablette) 5 keer per dag met 4-uurlikse intervale indien kreatinienopruiming > 25 ml / minuut is.

Pasiënte wat hemodialise ondergaan, moet hul gewone toepaslike dosis na elke dialise ontvang

Mondelike toegediende asiklovir by kinders jonger as 2 jaar is nog nie volledig bestudeer nie.

Die tablette moet per mond geneem word.

Indien u meer ADCO ACYCLOVIR 200 mg neem as wat u moes

In die geval van oordosering, raadpleeg u dokter of apteker. Indien nie een beskikbaar is nie, kontak die naaste hospitaal of gifhulpentrum.

Indien u vergeet om ADCO ACYCLOVIR 200 mg te neem

Neem dit sodra u onthou. Indien dit egter amper tyd is vir u volgende dosis, slaan die vergete dosis oor.

Moenie 'n dubbele dosis neem om op te maak vir 'n vergete dosis nie.

4. Moontlike newe-effekte

ADCO ACYCLOVIR 200 mg kan newe-effekte hê.

Nie alle newe-effekte wat vir ADCO ACYCLOVIR 200 mg aangemeld is, word by hierdie biljet ingesluit nie. Indien u algemene gesondheid verswak of indien u enige ongewenste

gevolge ervaar terwyl u ADCO ACYCLOVIR 200 mg neem, raadpleeg asseblief u gesondheidsorgverskaffer vir advies.

Indien enige van die volgende gebeur, staak die gebruik van ADCO ACYCLOVIR 200 mg en lig u dokter dadelik in, of gaan na die ongevalle-afdeling by u naaste hospitaal:

Gereelde newe-effekte sluit in:

- hoofpyn
- voel duiselig
- naarheid
- diarree
- braking
- maagpyn
- jeukerige vel (pruritus)
- veluitslag
- velreaksie na blootstelling aan lig (fotosensitiwiteit)
- galbulte (urtikaria)
- haarverlies
- voel moeg
- koors

Minder gereelde newe-effekte sluit in:

- effekte op sommige bloedtoetse
- anafilakse (lewensgevaarlike allergiese reaksie)
- angio-edeem (vinnige swelling onder die vel)
- agitاسie, verwarring, beweringheid, verswakte koördinasie, onduidelike spraak, hallusinasies, psigotiese simptome, stuipaanvalle, slaperigheid, breinsiektes, koma
- kortasem
- omkeerbare stygings in bilirubien en lewerverwante ensieme
- inflammasie van die lewer (hepatitis)
- vergeling van u vel en wit van u oë (geelsug)
- verhoogde bloedureum en kreatinien
- nierversaking, nierpyn

Indien u enige newe-effekte opmerk wat nie in hierdie biljet genoem word nie, stel asseblief u dokter of apteker in kennis.

Aanmelding van newe-effekte

Indien u newe-effekte ervaar, raadpleeg u dokter, apteker of verpleegkundige. U kan ook newe-effekte by SAHPRA aanmeld via die vorm "**6.04 Adverse Drug Reaction Reporting Form**", wat aanlyn gevind kan word onder SAHPRA se publikasies: <https://www.sahpra.org.za/Publications/Index/8>. Deur newe-effekte aan te meld, kan u help om meer inligting oor die veiligheid van ADCO ACYCLOVIR 200 mg te verskaf.

U kan dit ook by Adcock Ingram Limited aanmeld deur die volgende e-posadres te gebruik:
Adcock.AEReports@adcock.com.

5. Hoe om ADCO ACYCLOVIR 200 mg te bêre

- Bêre alle medisyne buite bereik van kinders.
- Bêre teen of benede 25 °C op 'n droë plek. Beskerm teen lig.
- Bêre in die oorspronklike verpakking.
- Moenie gebruik ná die vervaldatum wat op die etiket / kartonboksie / bottel vermeld word nie
- Neem alle ongebruikte medisyne terug na u apteker.
- Moenie ongebruikte medisyne in afvoerpype of rioolstelsels (bv. toilette) weggooi nie.

6. Inhoud van die verpakking en ander inligting

Wat ADCO ACYCLOVIR 200 mg bevat

Die aktiewe bestanddeel is asiklovir 200 mg.

Die ander bestanddele is laktose, kolloïdale silikondioksied, magnesiumstearaat, mikrokristallyne sellulose, voorafgelatiniseerde mieliestysel, natriumlourielsulfaat, natriumstyselglikolaat en steariensuur.

Hoe ADCO ACYCLOVIR lyk en die inhoud van die verpakking

ADCO ACYCLOVIR TABLETTE 200 mg is: Wit tablette, seskantig van vorm, breeklyn aan die een kant.

ADCO ACYCLOVIR 200 mg is beskikbaar in:

Verpakkingsgrootte van 25 tablette, verpak in 'n wit polipropileen-securitainer met 'n wit LDPE-sluiting.

Verpakkingsgrootte van 25 tablette verpak in stulpstroke met PVC-film en aluminiumfoelie. Nie alle verpakkingsgroottes word dalk bemark nie.

Houer van Sertifikaat van Registrasie

Adcock Ingram Limited

New Road 1,

Erand Gardens,

Midrand,

1685

Kliëntediens: 0860 ADCOCK / 232625

Hierdie biljet is mees onlangs hersien op

08 November 2024

Registrasienommer

28/20.2.8/0552

Datum van goedkeuring: 08 November 2024

PASIËNT INLIGTINGSBILJET

Botswana (S2): BOT1803283

Namibië (NS2): 04/20.2.8/1554

adcock ingram 

1229203 02/2025

Datum van goedkeuring: 08 November 2024