

## SCHEDULING STATUS

S3

### ADCO ALLOPURINOL 300 Tablets

#### Allopurinol

**Contains sugar (lactose):  
180 mg lactose per tablet**

### Read all of this leaflet carefully before you start taking ADCO ALLOPURINOL 300

- Keep this leaflet. You may need to read it again.
- If you have further questions, please ask your doctor, pharmacist, nurse or other health care provider.
- ADCO ALLOPURINOL 300 has been prescribed for you personally and you should not share your medicine with other people. It may harm them, even if their symptoms are the same as yours.

### What is in this leaflet

1. What ADCO ALLOPURINOL 300 is and what it is used for
2. What you need to know before you take ADCO ALLOPURINOL 300
3. How to take ADCO ALLOPURINOL 300
4. Possible side effects
5. How to store ADCO ALLOPURINOL 300
6. Contents of the pack and other information

#### 1. What ADCO ALLOPURINOL 300 is and what it is used for

ADCO ALLOPURINOL 300 is used for the treatment of gout and to prevent other conditions caused by hyperuricaemia (abnormally high levels of uric acid in the blood), such as kidney stones.

ADCO ALLOPURINOL 300 contains the active ingredient allopurinol and works by reducing the amount of uric acid in the blood.

#### 2. What you need to know before you take ADCO ALLOPURINOL 300

##### Do not take ADCO ALLOPURINOL 300:

- If you are allergic to allopurinol or to any of the ingredients in ADCO ALLOPURINOL 300 (listed in section 6).
- If you are pregnant or breastfeeding.
- If you have acute gout.

ADCO ALLOPURINOL 300 is not recommended in children, except those with abnormal cell growth (malignancy).

### Warnings and precautions

Take special care with ADCO ALLOPURINOL 300:

- if you experience any skin rash or allergic reaction. ADCO ALLOPURINOL 300 treatment should be stopped and contact your doctor for advice.

## PATIENT INFORMATION LEAFLET

---

- if you have an acute gout attack. ADCO ALLOPURINOL 300 treatment should not be started until the attack has subsided. Your doctor will advise on suitable treatment options to reduce inflammation during early stages of treatment.
- In children with Lesch-Nyhan syndrome (rare genetic disorder that affects mostly boys and causes them to produce too much uric acid). Increasing the daily fluid intake and taking urinary alkalinisers can minimize the formation of xanthine stones (kidney stones) in these children.
- if you suffer from or have a history of any urinary disorders, such as renal pelvic stones. Consult your doctor for advice.
- if you experience a fever or swelling under the skin
- if you have or ever had kidney or liver problems
- if you suffer from high blood pressure or a heart disorder and you are taking medicines to treat these conditions such as diuretics or ACE inhibitors.
- if you experience a fever, weight loss or fatigue, as these are symptoms of vasculitis (inflammation of blood vessels).
- if you experience seizures, difficulty breathing, sweating, fast heartbeat or sweating.

### **Other medicines and ADCO ALLOPURINOL 300**

Always tell your health care provider if you are taking any other medicine. (This includes all complementary or traditional medicines.)

Tell your doctor or pharmacist if you are currently taking:

- Mercaptopurine, azathioprine or any other anti-neoplastic medicines (used to treat certain types of cancers).
- Chlorpropamide and some sulfonylurea antidiabetics (used in the management of high blood sugar).
- Vidarabine (used to treat viral infections such as herpes simplex).
- Medicines that can increase uric acid concentration, such as aspirin or other salicylates used for pain and inflammation.
- Probenecid (used to treat chronic gout)
- Anticoagulants such as Warfarin (used to prevent blood clots)
- Phenytoin (used to treat seizures and epilepsy)
- Theophylline (used in asthma and other respiratory disorders)
- Some antibacterial medicines such as amoxicillin or ampicillin (used to treat bacterial infections)
- Ciclosporin (used in organ transplants)
- Didanosine (used in the treatment of HIV/AIDS)

### **ADCO ALLOPURINOL 300 with food and drink**

See section 3.

### **Pregnancy, breastfeeding and fertility**

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other health care provider for advice before taking this medicine.

### **Pregnancy**

You should not take ADCO ALLOPURINOL 300 if you are pregnant.

### **Breastfeeding**

You should not take ADCO ALLOPURINOL 300 if you are breastfeeding.

### **Fertility**

There is no information available on the effects of fertility.

### **Driving and using machines**

ADCO ALLOPURINOL 300 can cause vision tiredness, light-headedness and dizziness in some people. If this happens to you, avoid activities which require you to be alert; for example, driving a car and using machinery.

It is not always possible to predict to what extent ADCO ALLOPURINOL 300 may interfere with the daily activities of a patient. Patients should ensure that they do not engage in the above activities they are aware of the measure to which ADCO ALLOPURINOL 300 affects them.

### **ADCO ALLOPURINOL 300 contains lactose**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### **3. How to take ADCO ALLOPURINOL 300**

Do not share medicines prescribed for you with any other person.

Always take ADCO ALLOPURINOL 300 exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Doses up to and including 300 mg of ADCO ALLOPURINOL 300 may be taken once a day. Your doctor will decide on the correct dose, depending on your body's response and by monitoring your uric acid serum levels.

In severe conditions, doses of up to 600 mg may be required.

It is recommended that ADCO ALLOPURINOL 300 be taken after meals with water for better tolerance and to avoid nausea and vomiting.

### **Children**

The suggested initial dose is 8 mg/kg body mass daily.

You should take adequate amounts of fluid (at least 2 litres) during the day while on treatment with ADCO ALLOPURINOL 300.

The doses will be reduced for patients with kidney problems.

Your doctor will tell you how long your treatment with ADCO ALLOPURINOL 300 will last. Do not stop treatment early without consulting your health care provider first.

If you have the impression that the effect of ADCO ALLOPURINOL 300 is too strong or too weak, tell your doctor or pharmacist.

### **If you take more ADCO ALLOPURINOL 300 than you should**

## PATIENT INFORMATION LEAFLET

---

In the event of an overdose, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison centre.

Take the medicine pack with you. This is so the doctor knows what you have taken.

The most likely reaction would be gastro-intestinal intolerance.

Other symptoms of overdose include nausea, vomiting, diarrhoea and dizziness.

Treatment of overdose involves supportive measures such as administering sufficient fluids to facilitate excretion of ADCO ALLOPURINOL 300 from the body.

If necessary, haemodialysis may be used (a process whereby a machine filters and cleans the blood in the body).

### **If you forget to take ADCO ALLOPURINOL 300**

Do not take a double dose to make up for the forgotten individual doses.

### **If you stop taking ADCO ALLOPURINOL 300**

Talk to your doctor before you stop taking ADCO ALLOPURINOL 300.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

## **4. Possible side effects**

ADCO ALLOPURINOL 300 can have side effects.

Not all side effects reported for ADCO ALLOPURINOL 300 are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking ADCO ALLOPURINOL 300, please consult your health care provider for advice.

Tell your doctor immediately if you notice any of the following:

*Side effects occurring less frequently:*

- Furunculosis (pus-filled bumps under the skin caused by infection of the hair follicles)
- Aplastic anaemia (a condition that causes the bone marrow to stop producing new blood cells)
- Thrombocytopenia (a condition in which your body has too less platelets)
- Leukopenia (low levels of white blood cells) or Leukocytosis and Eosinophilia (high) levels of white blood cells
- Bone marrow depression (a condition in which the bone marrow cannot make enough blood cells)
- Haemolytic anaemia (excessive breakdown of red blood cells)
- Agranulocytosis (very low level of a type of white blood cell, granulocytes)
- Skin rash caused by an allergic reaction, Stevens-Johnson syndrome (life-threatening reaction with flu-like symptoms and painful rash affecting the skin, mouth, eyes and genitals), toxic epidermal necrolysis (skin disorder that causes blistering and peeling of skin), enlarged lymph nodes, flaky skin and angioimmunoblastic lymphadenopathy.
- Vasculitis (inflammation of blood vessels)
- Joint pain
- Drowsiness
- Diabetes
- High blood levels fat in the blood (Hyperlipidaemia)
- Depression

## PATIENT INFORMATION LEAFLET

---

- Dizziness
- Headache
- Ataxia (inability to coordinate muscle movements)
- Paraesthesia (sensations like numbness, tingling, pins and needles)
- Seizures
- Coma
- Paralysis
- Weakness, numbness and pain in the hands and feet from nerve damage
- Cataract (clouding of the lens)
- Visual disturbances
- Macular changes (an eye disease that causes vision loss)
- Vertigo (a spinning sensation)
- Angina (pains to the chest, jaw and back, brought on by physical effort and due to problems with the blood flow to the heart)
- Bradycardia (slow heart rate)
- Hypertension (high blood pressure)
- Nausea
- Vomiting
- Abdominal pain
- Diarrhoea
- Gastric irritation
- Taste disturbances
- Recurrent haematemesis (vomiting blood)
- Steatorrhoea (oily, smelly stools, which often float)
- Stomatitis (inflammation of the lining of the mouth)
- Changed bowel habit
- Liver damage and impaired liver function
- Inflammation of the liver
- Hair loss
- Angioedema (swelling under the skin surface due to an allergic reaction)
- Discoloured hair
- Muscle aches
- Kidney damage
- Haematuria (blood in the urine)
- Uraemia (high levels of urea in the blood)
- Gynaecomastia (abnormal breast enlargement)
- Male infertility
- Erectile dysfunction
- Fever and chills
- Malaise (feeling generally unwell)
- Oedema (fluid retention)
- Asthenia (weakness)
- Sleepiness

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

### Reporting of side effects

If you get side effects, talk to your doctor or pharmacist. You can also report side effects to SAHPRA via the Med Safety APP (Medsafety X SAHPRA) and eReporting platform (who-umc.org) found on SAHPRA's website. By reporting side effects, you can help provide more information on the safety of ADCO ALLOPURINOL 300.

### 5. How to store ADCO ALLOPURINOL 300

- Store all medicines out of reach of children.
- Store at or below 30 °C.
- Protect from light and moisture
- Return all unused medicine to your pharmacist.
- Do not dispose of unused medicine in drains and sewerage systems (e.g. toilets).

### 6. Contents of the pack and other information

#### What ADCO ALLOPURINOL 300 contains

The active substance is allopurinol.

Each tablet contains 300 mg allopurinol.

The other ingredients are lactose, maize starch, croscarmellose sodium, povidone K25.

#### What ADCO ALLOPURINOL 300 looks like and contents of the pack

Tablets.

White, bisected, biconvex tablets.

30 and 250 tablets packed in white polypropylene securitainers with LDPE (low density polyethylene) closures.

30 and 250 tablets packed in white HDPE (high density polyethylene) containers with HDPE closures.

30 and 1000 tablets packed in clear PVC film/ printed aluminium foil blister packs in a cardboard carton.

#### Holder of Certificate of Registration

Adcock Ingram Limited

1 New Road

Erand Gardens

Midrand, 1685

Customer Care: 0860 ADCOCK / 232625

#### The leaflet was last revised in

26 November 2024

#### Registration number

29/3.3/0229

## SKEDULERINGSTATUS

S3

**ADCO ALLOPURINOL 300 tablette**

**Allopurinol**

**Bevat suiker (laktose):**

**180 mg laktose per tablet**

**Lees hierdie hele voubiljet noukeurig deur voordat u ADCO ALLOPURINOL 300 begin neem**

- Hou hierdie voubiljet. U sal dit dalk weer moet lees.
- Indien u verdere vrae het, raadpleeg asseblief u dokter, apteker, verpleegkundige of ander gesondheidsorgverskaffer.
- ADCO ALLOPURINOL 300 is aan u persoonlik voorgeskryf en u moet nie u medisyne met ander mense deel nie. Dit kan hulle benadeel, selfs al ervaar hul simptome dieselfde as u.

### **Wat in hierdie voubiljet is**

1. Wat ADCO ALLOPURINOL 300 is en waarvoor dit gebruik word
2. Wat u moet weet voordat u ADCO ALLOPURINOL 300 neem
3. Hoe om ADCO ALLOPURINOL 300 te neem
4. Moontlike newe-effekte
5. Hoe om ADCO ALLOPURINOL 300 te bêre
6. Inhoud van die verpakking en ander inligting

#### **1. Wat ADCO ALLOPURINOL 300 is en waarvoor dit gebruik word**

ADCO ALLOPURINOL 300 word gebruik vir die behandeling van jig en om ander toestande te voorkom wat veroorsaak word deur hiperurisemie (abnormaal hoë vlakke van uriensuur in die bloed), soos nierstene.

ADCO ALLOPURINOL 300 bevat die aktiewe bestanddeel allopurinol en werk deur die hoeveelheid uriensuur in die bloed te verminder.

#### **2. Wat u moet weet voordat u ADCO ALLOPURINOL 300 neem**

##### **Moenie ADCO ALLOPURINOL 300 neem nie:**

- Indien u allergies is vir allopurinol of vir enige van die bestanddele in ADCO ALLOPURINOL 300 (gelys in afdeling 6).
- Indien u swanger is of borsvoed.
- Indien u akute jig het.

ADCO ALLOPURINOL 300 word nie aanbeveel by kinders nie, behalwe dié met abnormale selgroei (kwaadaardigheid).

### **Waarskuwings en voorsorgmaatreëls**

Wees veral versigtig met ADCO ALLOPURINOL 300:

- indien u enige veluitslag of allergiese reaksie ervaar. ADCO ALLOPURINOL 300 behandeling moet gestaak word en kontak u dokter vir advies.

## PASIËNT INLIGTINGSVOUBILJET

---

- indien u 'n akute jigaanval het. Behandeling met ADCO ALLOPURINOL 300 moet nie begin word voordat die aanval bedaar het nie. U dokter sal u adviseer oor geskikte behandelingsopsies om inflammasie tydens vroeë stadiums van behandeling te verminder.
- By kinders met Lesch-Nyhan-sindroom (seldsame genetiese afwyking wat meestal seuns affekteer en veroorsaak dat hulle te veel uriensuur produseer). Die verhoging van daaglikse vloeistofinname en die neem van urinêre alkaliseerders kan die vorming van xantien-stene (nierstene) by hierdie kinders verminder.
- indien u ly aan, of 'n geskiedenis het van enige urinêre afwykings, soos nierbekkenstene. Raadpleeg u dokter vir advies.
- indien u koors of swelling onder die vel ervaar
- indien u nier- of lewerprobleme het of ooit gehad het
- indien u aan hoë bloeddruk of 'n hartverstoring ly en u neem medisyne om hierdie toestande te behandel, soos diuretika of AOE-remmers.
- indien u koors, gewigsverlies of moegheid ervaar, aangesien dit simptome van vaskulitis (ontsteking van bloedvate) is.
- indien u stuipaanvalle, moeilike asemhaling, sweet, vinnige hartklop of sweet ervaar.

### Ander medisyne en ADCO ALLOPURINOL 300

Lig altyd u gesondheidsorgverskaffer in indien u enige ander medisyne neem. (Dit sluit alle komplementêre of tradisionele medisyne in.)

Lig u dokter of apteker in indien u tans neem:

- Merkaptopurien, azatioprien of enige ander anti-neoplastiese medisyne (wat gebruik word om sekere soorte kankers te behandel).
- Chloorpropamied en sommige sulfonyleurea-antidiabetika (gebruik in die beheer van hoë bloedsuiker).
- Vidarabien (word gebruik om virusinfeksies soos “herpes simplex” te behandel).
- Medisyne wat uriensuurkonsentrasie kan verhoog, soos aspirien of ander salisilate wat vir pyn en inflammasie gebruik word.
- Probenesied (word gebruik om chroniese jig te behandel)
- Antikoagulate soos Warfarien (gebruik om bloedklonte te voorkom)
- Fenitoïen (word gebruik om stuipaanvalle en epilepsie te behandel)
- Teofillien (gebruik in asma en ander respiratoriese afwykings)
- Sommige antibakteriese medisyne soos amoksisillien of ampisillien (word gebruik om bakteriële infeksies te behandel)
- Siklosporien (gebruik in orgaanoorplantings)
- Didanosien (gebruik in die behandeling van MIV/VIGS)

### ADCO ALLOPURINOL 300 saam met kos en drank

Sien afdeling 3.

### Swangerskap, borsvoeding en vrugbaarheid

Indien u swanger is of borsvoed, dink u kan swanger wees of van plan is om 'n baba te hê, raadpleeg asseblief u dokter, apteker of ander gesondheidsorgverskaffer vir advies voordat u hierdie medisyne neem.

### Swangerskap



U moet nie ADCO ALLOPURINOL 300 neem indien u swanger is nie.

### **Borsvoeding**

U moet nie ADCO ALLOPURINOL 300 neem indien u borsvoed nie.

### **Vrugbaarheid**

Daar is geen inligting beskikbaar oor die gevolge op vrugbaarheid nie.

### **Bestuur en gebruik van masjinerie**

ADCO ALLOPURINOL 300 kan visie-moegheid, lighoofdigheid en duiseligheid by sommige mense veroorsaak. Indien dit met u gebeur, vermy aktiwiteite wat vereis dat u waaksaam moet wees; byvoorbeeld om 'n motor te bestuur en masjinerie te gebruik.

Dit is nie altyd moontlik om te voorspel tot watter mate ADCO ALLOPURINOL 300 die daaglikse aktiwiteite van 'n pasiënt kan beïnvloed nie. Pasiënte moet verseker dat hulle nie bogenoemde aktiwiteite verrig totdat hulle bewus is van die mate waartoe ADCO ALLOPURINOL 300 hulle beïnvloed nie.

### **ADCO ALLOPURINOL 300 bevat laktose**

Indien u deur u dokter ingelig is dat u 'n onverdraagsaamheid teenoor sommige suikers het, kontak u dokter voordat u hierdie medisyne neem.

### **3. Hoe om ADCO ALLOPURINOL 300 te neem**

Moenie medisyne wat aan u voorgeskryf is, met enige ander persoon deel nie.

Neem altyd ADCO ALLOPURINOL 300 presies soos u dokter of apteker u ingelig het. Raadpleeg u dokter of apteker indien u nie seker is nie.

Dosisse tot en met 300 mg ADCO ALLOPURINOL 300 kan een keer per dag geneem word. U dokter sal besluit oor die korrekte dosis, afhangende van u liggaam se reaksie, en deur u uriensuurserumvlakke te monitor.

In ernstige toestande kan dosisse van tot 600 mg benodig word.

Dit word aanbeveel dat ADCO ALLOPURINOL 300 na etes met water geneem word vir beter verdraagsaamheid en om naarheid en braking te voorkom.

### **Kinders**

Die voorgestelde aanvanklike dosis is 8 mg/kg liggaamsmassa eenmaal per dag.

U moet voldoende hoeveelhede vloeistof (ten minste 2 liter) gedurende die dag neem terwyl u met ADCO ALLOPURINOL 300 behandel word.

Die dosisse sal verminder word vir pasiënte met nierprobleme.

U dokter sal u inlig hoe lank u behandeling met ADCO ALLOPURINOL 300 sal duur. Moenie die behandeling voortydig stop sonder om eers u gesondheidsorgverskaffer te raadpleeg nie.

Indien u die indruk het dat die effek van ADCO ALLOPURINOL 300 te sterk of te swak is, lig u

dokter of apteker in.

### **Indien u meer ADCO ALLOPURINOL 300 neem indien wat u moes**

In die geval van 'n oordosis, raadpleeg u dokter of apteker. Indien nie een beskikbaar is nie, kontak die naaste hospitaal of gifhulpentrum.

Neem die medisyne-verpakking saam. Dit is sodat die dokter weet wat u geneem het.

Die mees waarskynlike reaksie sou gastro-intestinale onverdraagsaamheid wees.

Ander simptome van oordosis sluit in naarheid, braking, diarree en duiseligheid.

Behandeling van oordosis behels ondersteunende maatreëls soos die toediening van voldoende vloeistowwe om die uitskeiding van ADCO ALLOPURINOL 300 uit die liggaam aan te help.

Indien nodig, kan hemodialise gebruik word ('n proses waardeur 'n masjien die bloed in die liggaam filtreer en skoonmaak).

### **Indien u vergeet om ADCO ALLOPURINOL 300 te neem**

Moenie 'n dubbele dosis neem om op te maak vir die vergete individuele dosis nie.

### **Indien u die gebruik van ADCO ALLOPURINOL 300 staak**

Raadpleeg u dokter voordat u ophou om ADCO ALLOPURINOL 300 te neem.

Indien u enige verdere vrae het oor die gebruik van hierdie medisyne, vra u dokter, apteker of verpleegkundige.

## **4. Moontlike nuwe-effekte**

ADCO ALLOPURINOL 300 kan nuwe-effekte hê.

Nie alle nuwe-effekte wat vir ADCO ALLOPURINOL 300 aangemeld is, word in hierdie voubiljet ingesluit nie. Indien u algemene gesondheid vererger of u enige ongewenste gevolge ervaar terwyl u ADCO ALLOPURINOL 300 neem, raadpleeg asseblief u gesondheidsorgverskaffer vir advies.

Lig u dokter dadelik in indien u enige van die volgende opmerk:

*Nuwe-effekte wat minder algemeen voorkom:*

- Furunkulose (etter-ge vulde knoppe onder die vel wat veroorsaak word deur infeksie van die haarfollikels)
- Aplastiese anemie ('n toestand wat veroorsaak dat die beenmurg ophou om nuwe bloedselle te produseer)
- Trombositopenie ('n toestand waarin u liggaam te min bloedplaatjies het)
- Leukopenie (lae vlakke van witbloedselle) of leukositose en eosinofilie (hoë) vlakke van witbloedselle
- Beenmurgdepressie ('n toestand waar die beenmurg nie genoeg bloedselle kan maak nie)
- Hemolitiese anemie (oormatige afbraak van rooibloedselle)
- Agranulositose (baie lae vlak van 'n tipe witbloedsel, granulose)
- Veluitslag veroorsaak deur 'n allergiese reaksie, Stevens-Johnson-sindroom (lewensgevaarlike reaksie met griepagtige simptome en pynlike uitslag wat die vel, mond, oë en geslagsdele aantast), toksiese epidermale nekrolise (velafwyking wat blase en afskilfering van die vel

## PASIËNT INLIGTINGSVOUBILJET

---

veroorzaak), vergrote limfknope, skilferige vel en angio-immunoblastiese limfadenopatie.

- Vaskulitis (ontsteking van bloedvate)
- Gewrigspyn
- Lomerigheid
- Diabetes
- Hoë bloedvlakke van vet in die bloed (hiperlipidemie)
- Depressie
- Duiseligheid
- Hoofpyn
- Ataksie (onvermoë om spierbewegings te koördineer)
- Parestesie (sensasies soos gevoelloosheid, tinteling, naalde-en-spelde)
- Stuipaanvalle
- Koma
- Verlamming
- Swakheid, gevoelloosheid en pyn in die hande en voete as gevolg van senuweeskade
- Katarak (vertroebeling van die lens)
- Visuele verstourings
- Makulêre veranderinge ('n oogsiekte wat visieverlies veroorsaak)
- Vertigo ('n tollende sensasie)
- Angina (pyn in die bors, kakebeen en rug, veroorsaak deur fisieke inspanning en as gevolg van probleme met die bloedvloei na die hart)
- Bradikardie (stadige hartklop)
- Hipertensie (hoë bloeddruk)
- Naarheid
- Braking
- Buikpyn
- Diarree
- Maag irritasie
- Smaak-verstourings
- Herhalende hematemes (braking van bloed)
- Steatoree (olierige, onwelriekende ontlasting, wat dikwels dryf)
- Stomatitis (ontsteking van die voering van die mond)
- Veranderde ontlastingsroetine
- Lewerskade en verswakte lewerfunksie
- Inflammasie van die lewer
- Haarverlies
- Angio-edeem (swelling onder die veloppervlak as gevolg van 'n allergiese reaksie)
- Verkleurde hare
- Spierpyn
- Nierskade
- Hematurie (bloed in die urien)
- Uremie (hoë vlakke van ureum in die bloed)
- Ginekomastie (abnormale borsvergroting)
- Manlike onvrugbaarheid
- Erektiele disfunksie
- Koors en kouekoors

## PASIËNT INLIGTINGSVOUBILJET

---

- Malaise (voel oor die algemeen onwel)
- Edeem (vloeistofretensie)
- Astenie (swakheid)
- Slaperigheid

Indien u enige newe-effekte opmerk wat nie in hierdie voubiljet genoem word nie, stel asseblief u dokter of apteker in kennis.

### **Aanmelding van newe-effekte**

Indien u newe-effekte kervaar, raadpleeg u dokter of apteker. U kan ook newe-effekte by SAHPRA aanmeld via die Med Safety Toep (Medsafety X SAHPRA) en elektroniese aanmeldingsplatform (who-umc.org) wat op SAHPRA se webwerf gevind kan word. Deur newe-effekte aan te meld, kan u help om meer inligting oor die veiligheid van ADCO ALLOPURINOL 300 te verskaf.

### **5. Hoe om ADCO ALLOPURINOL 300 te bêre**

- Bêre alle medisyne buite bereik van kinders.
- Bêre teen of benede 30 °C.
- Beskerm teen lig en vog
- Neem alle ongebruikte medisyne terug na u apteker.
- Moenie ongebruikte medisyne in afvoerpype en rioolstelsels (bv. toilette) weggooi nie.

### **6. Inhoud van die verpakking en ander inligting**

#### **Wat ADCO ALLOPURINOL 300 bevat**

Die aktiewe bestanddeel is allopurinol.

Elke tablet bevat 300 mg allopurinol.

Die ander bestanddele is laktose, meliestysel, natriumkruiskarmellose, povidoon K25.

#### **Hoe lyk ADCO ALLOPURINOL 300 en die inhoud van die verpakking**

Tablette.

Wit, gekeefte, bikonvekse tablette.

30 en 250 tablette verpak in wit polipropileen veiligheidshouers met LDPE (lae digtheid poliëtileen) sluitings.

30 en 250 tablette verpak in wit HDPE (hoëdigtheid poliëtileen) houers met HDPE-sluitings.

30 en 1000 tablette verpak in deursigtige PVC-film/ gedrukte aluminiumfoelie stulpstrook-verpakkings in 'n kartonboksie.

#### **Houer van Registrasiesertifikaat**

Adcock Ingram Limited

New Road 1

Erand Gardens

Midrand, 1685

Kliëntediens: 0860 ADCOCK / 232625

## PASIËNT INLIGTINGSVOUBILJET

---

**Die voubiljet is mees onlangs hersien op**

26 November 2024

**Registrasienuommer**

29/3.3/0229

**adcock ingram** 

1229002 12/2024

**Datum van goedkeuring:** 26 November 2024